


EE212 Mathematical Foundations for Machine Learning and Data Science
TUTORIAL 3

2020 Mid-Exam Review
Tutorial 3-1

In each of the plot given below, we have plotted two vectors $x, y \in \mathbf{R}^n$ such that each point represents (x_i, y_i) for $i = 1, 2, \dots, n$. In each case, determine whether the correlation coefficient ρ of two vectors is positive ($\rho > 0.5$), negative ($\rho < -0.5$) or near zero (say $|\rho| < 0.3$).



Tutorial 3-2

What are the properties of a norm function? Determine 1-norm, Euclidean norm and ∞ -norm of a vector $x = (7, -1, 2, -1, 3)$.

Tutorial 3-3

Let $A \in \mathbf{R}^{m \times n}$ represent a contingency matrix of m cabinet members who hold n residencies (or citizenships)

$$A_{ij} = \begin{cases} 1 & \text{member } i \text{ holds residency } j \\ 0 & \text{member } i \text{ does not hold residency } j, \end{cases}$$

where we note that a member can hold any number of residencies.

- (a) Give a simple formula for the n -vector g , where g_i is the total number of members holding residency j .

Give simple English description of the following expressions or quantities.

- (b) 2-nd column of A .
(c) $(AA^T)_{ij}$
(d) $(A^T A)_{ij}$

Tutorial 3-4

Which of following matrices are left-invertible? Provide justification to support your answer.

(a) $A = \begin{bmatrix} -1 & 2 \\ 3 & -6 \\ 2 & -1 \end{bmatrix}$

(b) $A = \begin{bmatrix} -1 & 3 & 2 \\ 2 & -6 & -1 \end{bmatrix}$

(c) $A = \begin{bmatrix} -9 & 0 & 7 \\ 4 & 0 & -5 \\ -1 & 0 & 6 \end{bmatrix}$

(d) $A = \begin{bmatrix} B \\ C \end{bmatrix}$, where $B \in \mathbf{R}^{n \times n}$ is a diagonal matrix with non-zero diagonal entries and $C \in \mathbf{R}^{m \times n}$ can be any matrix.

(e) $A = ab^T$, where $a, b \in \mathbf{R}^n$ for $n > 1$.

Tutorial 3-5

Use Gram-Schmidt process to determine the QR factorization of the matrix A given by

$$A = \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{\sqrt{2}} & 1 \\ 0 & 1 \end{bmatrix}.$$

Tutorial 3-6

Consider a matrix A given by

$$A = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}.$$

- (a) Give simple description of the relationship between x and Ax .
- (b) Determine A^5 .

Tutorial 3-7

(Independent parts)

- (a) Consider two linearly independent vectors $a_1, a_2 \in \mathbf{R}^3$. Consider a set W that contains all points of the form $\beta_1 a_1 + \beta_2 a_2 + 3$. Is W a subspace? If yes, what is the dimension of the subspace?
- (b) Consider a vector space $V = \mathbf{R}^m$ and the span of the vectors a_1, a_2, \dots, a_n of the vector space V . What is the maximum possible dimension of the span of the vectors a_1, a_2, \dots, a_n of the vector space?

Tutorial 3-8

Let $A \in \mathbf{R}^{3 \times 2}$ be a matrix given by

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 2 \\ 2 & 1 \end{bmatrix},$$

with singular-value decomposition given as $A = U\Sigma V^T$. The eigenvalue decomposition of $B = A^T A$ has 1 and 17 as its two non-zero eigenvalues and one of the two normalised eigenvectors is

$$q = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \end{bmatrix}^T$$

- (a) Determine the matrix V .
(Hint: You only need to use the given information above and exploit the fact that V is an orthogonal matrix.)
- (b) Determine the singular values of A , that is, the matrix Σ .
(Hint: Relate the EVD of B with the SVD of A)
- (c) Determine the determinant of the matrix B .